

IBUPROFEN (MOTRIN or ADVIL) DOSING

WEIGHT IN POUNDS (lbs) and APPROXIMATE AGE	**Dose in milligrams (mg)**	<u>Infant Concentrated Drops</u> 50mg/1.25ml	<u>Children's Liquid Suspension</u> 100mg/5ml	<u>Children's Chewable Tablets</u> 50 mg each	<u>Junior Strength Chewable or Non-chewable Tablets</u> 100 mg each	<u>Adult Tablet</u> 200 mg tablet
Under 6 months	Ibuprofen is Not Recommended for Infants less than 6 months old. If a fever (temperature greater than or equal to 100.4 F taken rectally) occurs in a child less than 3 months old please notify Pediatrician immediately.					
12-17 lbs 6-11 months	50mg	1.25ml	1/2 teaspoon	----	----	----
18-23 lbs 12-23 months	75 mg	1.875ml	3/4 teaspoon	----	----	----
24-35 lbs 2-3 years	100 mg	----	1 teaspoon	2 tablets	----	----
36-47 lbs 4-5 years	150 mg	----	1 1/2 teaspoon	3 tablets	----	----
48-59 lbs 6-8 years	200 mg	----	2 teaspoons	4 tablets	2 tablets	----
60-71 lbs 9-10 years	250 mg	----	2 1/2 teaspoons	5 tablets	2 1/2 tablets	----
72-95 lbs 11 years	300 mg	----	3 teaspoons	6 tablets	3 tablets	----
96 lbs and over 12 years old	400 mg	----	-----	----	4 tablets	2 tablets

Use of weight to select dose is preferred. If weight is not available, then use age.

May repeat dose after 6 hours. Do not exceed 4 doses in 24 hours.

If a child appears severely ill, giving ibuprofen is NOT a substitute for medical advice. **Please notify Pediatrician if you have concerns.**